



HAL
open science

**Conformational changes in amyloid-beta (12-28)
alloforms studied using action-FRET, IMS and
molecular dynamics simulations (vol 6, pg 5040, 2015)**

Steven Daly, Alexander Kulesza, Frederic Poussigue, Anne-Laure Simon,
Chang Min Choi, Geoffrey Knight, Fabien Chiro, Luke Macaleese, Rodolphe
Antoine, Philippe Dugourd

► **To cite this version:**

Steven Daly, Alexander Kulesza, Frederic Poussigue, Anne-Laure Simon, Chang Min Choi, et al..
Conformational changes in amyloid-beta (12-28) alloforms studied using action-FRET, IMS and
molecular dynamics simulations (vol 6, pg 5040, 2015). *Chemical Science*, 2016, 7, pp.1609-1610.
10.1039/c5sc90069g . hal-02303924

HAL Id: hal-02303924

<https://univ-lyon1.hal.science/hal-02303924>

Submitted on 25 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



CrossMark
click for updates

Cite this: *Chem. Sci.*, 2016, 7, 1609

Correction: Conformational changes in amyloid-beta (12–28) alloforms studied using action-FRET, IMS and molecular dynamics simulations

Steven Daly,^{ab} Alexander Kulesza,^{ab} Frederic Poussigue,^{ac} Anne-Laure Simon,^{ab} Chang Min Choi,^{ab} Geoffrey Knight,^{ab} Fabien Chiroi,^{ac} Luke MacAleese,^{ab} Rodolphe Antoine^{ab} and Philippe Dugourd^{*ab}

DOI: 10.1039/c5sc90069g

www.rsc.org/chemicalscience

Correction for 'Conformational changes in amyloid-beta (12–28) alloforms studied using action-FRET, IMS and molecular dynamics simulations' by Steven Daly *et al.*, *Chem. Sci.*, 2015, 6, 5040–5047.

There has been a consistent error in the name of the PDB code which was provided in the paper. The code given was 1LFM, but should have read 2LFM. There are 3 instances of this in the manuscript and one in the ESI.

(1) The inset of Fig. 2 and the figure caption should appear as:




Fig. 2 Representative structures simulated at 292 K of the dominant conformational family of the different charge states of the wild (top) and F19P (bottom) alloforms of A β_{12-28} . Here, the donor chromophore (grafted to the C-terminal residue) is shown in red, the acceptor chromophore in blue, and the peptide backbone in green. The inset on the top panel shows the NMR structure (pdb database file 2LFM) for the full amyloid beta protein, with the 12–28 region highlighted in green. The corresponding figure of the ensembles with the alternative chromophore grafting location is shown in Fig. S2.

^aUniversité de Lyon, F-69622, Lyon, France. E-mail: philippe.dugourd@univ-lyon1.fr

^bCNRS et Université Lyon 1, UMR5306, Institut Lumière Matière, France

^cCNRS et Université Lyon 1, UMR 5280, Institut des Sciences Analytiques, France



(2) On page 5044 (manuscript page 5), right hand column, line 12. The PDB code should read 2LFM rather than 1LFM.

(3) On page 5044, in the caption of Fig. 3. The caption should read:

Ramachandran plots for the hydrophobic core region (residues 17–21) of the different charge states of the wild (top) and F19P (bottom) alloforms of $A\beta_{12-28}$ (donor chromophore is grafted to the C-terminal residue). Black squares correspond to the 5 dihedral angle (Φ, Ψ) pairs of residues 17–21 for all the structures computed at 292 K. Red dots indicate the corresponding dihedral angles of the partially folded solution structure from the pdb file 2LFM. Blue dots indicate the dihedral angles of the A-chains in the pdb file 2BEG. The corresponding figure of the ensembles with the alternative chromophore grafting location is given in Fig. S3.†

(4) In the ESI, the caption of Fig. S3 should read:

Fig. S3. Ramachandran plots for the hydrophobic core region (residues 17–21) of the different charge states of the wild (top) and F19P (bottom) alloforms of $A\beta_{12-28}$ (donor chromophore is grafted to the N-terminal residue). Black squares correspond to the 5 dihedral angle (Φ, Ψ) pairs of residues 17–21 for all the structures computed at 292 K. Red dots indicate the corresponding dihedral angles of the partially folded solution structure from the pdb file 2LFM. Blue dots indicate the dihedral angles of the A-chains in the pdb file 2BEG.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

