Bottom-up processes control benthic macroinvertebrate communities and food web structure of fishless artificial wetlands

Florian Mermillod-Blondin, Pierre Marmonier, Mélissa Tenaille, Damien Lemoine, Michel Lafont, Ross Vander Vorste, Laurent Simon, Laurence Volatier

► To cite this version:

Florian Mermillod-Blondin, Pierre Marmonier, Mélissa Tenaille, Damien Lemoine, Michel Lafont, et al.. Bottom-up processes control benthic macroinvertebrate communities and food web structure of fishless artificial wetlands. *Aquatic Ecology*, 2020, 54, pp.575-589. 10.1007/s10452-020-09760-2 . hal-02515841

HAL Id: hal-02515841

<https://univ-lyon1.hal.science/hal-02515841>

Submitted on 10 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

TITLE: Bottom-up processes control benthic macro-invertebrate communities and food web structure of fishless artificial wetlands

Authors: Florian Mermillod-Blondin F.^{1*}, Pierre Marmonier¹, Mélissa Tenaille¹, Damien G. Lemoine¹, Michel Lafont¹, Ross Vander Vorste², Laurent Simon¹, Laurence Volatier¹

¹ UMR 5023 LEHNA, Université de Lyon, CNRS, Université Claude Bernard Lyon 1, ENTPE, 6 rue Raphaël Dubois, 69622 Villeurbanne, France.

² Department of Biology, University of Wisconsin, WI 54601, La Crosse, USA.

* Corresponding author: Florian Mermillod-Blondin

UMR 5023 LEHNA, Université de Lyon, CNRS, Université Claude Bernard Lyon 1, ENTPE, 6 rue Raphaël Dubois, 69622 Villeurbanne.

E-mail: florian.mermillod-blondin@univ-lyon1.fr

Phone number: (33) 4 72 43 13 64

Fax number: (33) 4 72 43 11 41

Abstract

In freshwater environments, the relative contributions of top-down and bottom-up effects on invertebrate communities in relation to productivity are largely ecosystem dependent. Artificial wetlands are increasingly developed to compensate for the loss of natural wetlands however their trophic processes remain poorly studied. The present study aimed to evaluate the respective contributions of bottom-up and top-down processes in structuring benthic food webs of three artificial wetlands with varying levels of benthic primary productivity. We found that phototrophic-based food webs in our artificial wetlands were controlled from the bottom-up by primary productivity and algal biomass developing at the water-sediment interface. No significant top-down-control of herbivore species by invertebrate predators was detected even in the wetland with the highest productivity. Increased richness of invertebrate grazers and scrapers with benthic primary productivity and algal biomass might have dampened the trophic cascade from predators to primary producers. In contrast with the phototrophic-based food web, analyses performed on the detritus-based food web showed that deposit-feeder invertebrate abundance was not correlated with the quantity of organic matter in sediments, suggesting a weak bottom-up effect of sedimentary organic matter content on deposit-feeders. More surprisingly, deposit-feeders, especially aquatic oligochaetes, seemed to influence the detritus-based food webs by stimulating organic matter processing and bacterial growth through bioturbation. The present study highlighted the occurrence of contrasting trophic processes between phototrophic-based and detritus-based food webs which can have implications on ecosystem functions, such as nutrient cycling and energy fluxes.

Keywords: functional feeding groups, benthic food webs, invertebrates, benthic biofilm, algae, bacteria, trophic relationships

Introduction

The relative importance of bottom-up and top-down controls on the structure of communities has long been an intense debate (Oksanen et al. 1981; Hunter and Price 1992; Polis and Strong 1996; Gruner et al. 2008). The exploitation ecosystem hypothesis (EEH) predicts that resource availability and predation control densities within adjacent trophic levels (Oksanen et al. 1981). Accordingly, increasing primary productivity lengthens food webs by boosting the establishment of predators, that will, in turn, have a greater impact on lower trophic levels. In nutrient-poor ecosystems, biomass of basal species is severely constrained by resource availability and is too low to support a significant consumer biomass (Menge 2000). In more nutrient-rich ecosystems, productivity of basal species increases with elevated resources but their biomass is kept constant by predation as consumers become more abundant (Steiner 2001). Although it has been shown that the predictions of EEH are modulated by biotic (e.g., diversity of the food webs) and abiotic (e.g., habitat heterogeneity, disturbance) factors (Hunter and Price 1992; Leibold et al. 1997; Menge et al. 2002; Foulquier et al. 2011), it provides a robust theoretical framework to evaluate the roles of bottom-up and top-down controls on ecosystem structure (e.g., Shurin et al. 2002; Aunapu et al. 2008; Lezama et al. 2014; Du et al. 2015; Hoset et al. 2017).

In wetlands, a historical view that bottom-up controls have greater relative importance than top-down controls in regulating ecosystem structure and function (e.g., Odum and Smalley 1959) has been reconsidered more recently (Sala et al. 2008; Moore 2018). Indeed, top predators like fish or invertebrates can significantly control the abundance and the assemblage of their prey (Diehl 1992; Magnusson and Williams 2009; Cobbaert et al. 2010; Laske et al. 2017) and top-down control of grazers on primary producers play a key role on wetland functioning and stability (e.g., limitation of algal blooms; Hillebrand 2009; Bertness et al. 2014). In this context, artificial wetlands, which

are increasingly developed to compensate for the loss of natural wetlands (Davidson 2014), could be useful tools for determining the respective roles of bottom-up and top-down forces on community structures because they allow for the selection of systems with distinct productivities. Moreover, these artificial systems may be managed for controlling the water flux of organisms, reducing the heterogeneity in biological colonization among wetlands (De Meester et al. 2005). We tested the predictions of EEH in shallow (water depth < 1.5 m) artificial wetlands having area of less than 2 ha and benthic habitat characterized by dense biofilms of phototrophic and heterotrophic micro-organisms (Gette-Bouvarot et al. 2014). To test the respective influences of bottom-up and top-down controls on benthic invertebrate communities, we selected three fishless shallow artificial wetlands characterized with contrasting primary productivities (estimated from net photosynthesis of benthic biofilms) associated with differences in sediment composition (proportion of clay and silt) and algal biomass at the water-sediment interface. We expected components of the biofilms to play a key role as food sources for the benthic food web creating a bottom-up effect. However, in accordance with the EEH we hypothesized the relative importance of bottom-up and top-down effects to shift from a dominance of bottom-up controls to a dominance of top-down controls with increasing primary productivity. To evaluate the response of food webs, we classified benthic invertebrate communities into trophic functional groups by including both the vertical (linear trophic chain from basal resources to predators through consumers) and horizontal dimensions of food webs (species diversity at each trophic level, following Srivastava and Bell (2009)). Under a bottom-up control scenario, we predicted a positive relationship between the abundance of grazers and scrapers and primary productivity and algal biomass at the water-sediment interface, a positive relationship between deposit-feeders (bacterivores) and bacterial abundance and organic matter (% of total organic carbon and total nitrogen) in sediment, and a positive relationship between macroinvertebrate predator abundance and the abundance of preys at

lower trophic levels. Alternatively, under a top-down control scenario, we predicted no positive relationships between consumers and their food sources.

Material and Methods

Study site

Our experiment took place in a pumping well field of « Crépieux-Charmy » (3.75 km²) located in the Rhône River floodplain (France), which provides drinking water for the metropolitan area of Lyon (1,280,000 inhabitants supplied). Artificial wetlands (surface area = 0.85-1.5 ha) have been built to recharge aquifer with river water. All wetlands were built by excavation in natural soils in areas with reduced riparian vegetation and were partially filled with a layer of calibrated sand about 20 cm thick and characterized by a low organic matter content (total organic carbon < 0.1% of sediment dry mass). Their main water supply is pumped in a channel (Vieux Rhône) of the Rhône River using centrifugal pumps (rotation: 1000 rounds per minute) that markedly limit the transport of top-predator fishes from Rhône River to wetlands. Despite similar designs, wetlands evolved over time due to the accumulation of fine particles (< 63 µm) at the water-sediment interface, influencing the availability of the benthic habitat for biofilms. Indeed, presence of fine sediments on a coarse and poorly organic sediment ameliorate the physical attachment and the nutrient availability for benthic biofilms (Sundbäck et al. 2003; van de Koppel et al. 2001; Watermann et al. 1999). For the present study, we selected three wetlands (called A, B, C) that were characterized by varying amounts of fine particles (proportions determined by laser diffractometry following Estragnat et al. 2018) at the water-sediment interface (Fig. 1A, one-way ANOVAs, comparison of fine sediment proportion among wetlands, $F_{2,15} = 29.6$, $p < 0.001$), leading to contrasted benthic

habitats for benthic biofilms. Following the decreasing gradient of fine particle amount from wetland A to wetland C, algal biomass and net photosynthesis activity (see below for details on measurements) of the biofilm decreased from wetland A with a mean chlorophyll a content of 2.85 $\mu\text{g per cm}^2$ and positive net production to wetland C with a mean chlorophyll a content of 1.55 $\mu\text{g per cm}^2$ and negative net production as photosynthesis did not compensate for respiration (Fig. 1BC, one-way ANOVAs, comparisons of algal biomass and net photosynthesis among wetlands, $F_{2,15} = 26.6$, $p < 0.001$ for algal biomass, $F_{2,15} = 23.6$, $p < 0.001$ for net photosynthesis). These conditions allowed us to evaluate the trophic relationships between primary productivity, benthic food sources, and invertebrate communities.

Environmental conditions before sampling date

The three studied wetlands had the same water supply and water column depth during a period of 6 weeks before sampling date (on 22nd of October 2014), minimizing the potential influence of hydrological conditions on community structure. During this period, we also verified that the three wetlands were characterized by comparable physico-chemical conditions in the water column. *In situ* monitoring of water temperatures showed similar variations (between 14.7 and 24.6 °C) in the three artificial wetlands. Weekly measurements of specific conductivity, pH, and dissolved oxygen (DO) concentrations performed with a HQ40D multiparameter (HACH) also indicated comparable values among artificial wetlands with mean ($\pm$ standard deviation) values of 337 (± 13) $\mu\text{S/cm}$, 8.3 (± 0.3), and 9.4 (± 0.5) mg/L for specific conductivity, pH and DO, respectively. Nutrient concentrations were also measured weekly in the water column of the wetlands following Gette-Bouvarot et al. (2014). Comparable dissolved nutrient concentrations and fluctuations were found in the three wetlands with concentrations of N-NO_3^- , N-NH_4^+ and P-PO_4^{3-} of 0.99 ± 0.21 (mean $\pm$ SD) mg/L, 32.78 ± 20.13 $\mu\text{g/L}$ and 12.29 ± 9.27 $\mu\text{g/L}$, respectively.

Sampling design

Biofilm and invertebrate samples were collected on the same day (October 22, 2014) to obtain a data set linking benthic fauna with characteristics of the sedimentary biofilm. Analyses on benthic samples were performed to determine biofilm characteristics (fluorescence measurements of benthic algal biomass and composition, bacterial abundance, total organic carbon and total nitrogen of the biofilm) and benthic communities of invertebrates. To facilitate the collection of benthic samples, the water pumping into the study wetlands was stopped 2 days before the sampling date to have between 30 and 40 cm of water level in all wetlands. In each wetland, we randomly selected 6 replicate zones. For biofilm analyses, the top layer (0-1 cm) of sediments was cored in each zone using cut syringes (internal diameter = 15 mm). Fifteen samples were collected, mixed and homogenized per zone to obtain representative sediment samples. For net photosynthetic activity of the biofilm, analyses were performed on one intact core collected per zone. Benthic invertebrates were sampled in each zone using a 200- μ m-mesh surber net following core collections for sediment and biofilm analyses. The top sediment layer was collected on a surface of 150 cm² per zone and preserved with 96% ethanol. All samples were then stored at ~10°C during transport to the laboratory within 4 hours.

Chemical and biological analyses on sediments

The biomass of green algae, diatoms and cyanobacteria at the water-sediment interface were estimated by *in situ* measurements of chlorophyll-*a* concentration and were expressed as chlorophyll-*a* equivalent per unit of surface. These measurements were performed with a benthic fluorimetric probe (BenthoTorch, bbe Moldaenke GmbH) (see Harris and Graham 2015 for details).

The bacterial cell numbers were evaluated on sediment previously fixed with formaldehyde (final concentration 2 % v/v) by direct microscopy counts after incubation of the sediment with the Syto9 green fluorescent nucleic acid stain (Invitrogen). Details of the protocol were given in Gette-Bouvarot et al. (2014). The total bacterial abundance (cells/g of dry sediment) was then determined.

Total organic carbon (TOC) and total nitrogen (TN) were measured as estimators of the sedimentary biofilm biomass. Following Mermillod-Blondin et al. (2015), TOC and TN were determined by high-temperature combustion of *in situ* pre-acidified (HCl 2N, to remove inorganic carbon) dry samples (60°C, 48h) and subsequent measurements of CO₂ and N₂ by thermal conductometry using an elemental analyzer (FlashEA 1112, Thermo Electron Corporation).

Net photosynthetic activity (NP) was determined by measuring oxygen production at 15 °C (close to field temperature at sampling date) on intact biofilms (4 cm² collected with a cut syringe, see above) using 4 mL respiration chambers fitted with micro-probes (Microrespiration system, Unisense, see Gette-Bouvarot et al. 2015 for details). Photosynthesis measurements were performed with an irradiance of 150 µmol.m⁻².s⁻¹, close to maximal photosynthetically active radiations (PAR) measured at midday with a LI-193 Spherical Quantum Sensor (PAR varying between 20 and 130 µmol m⁻² s⁻¹ depending on meteorological conditions). For measurements, total dissolved inorganic carbon concentration was adjusted to 10 mM with KHCO₃ to ensure sufficient supply of inorganic carbon during the incubation and pH was maintained at 6.5 by addition of a citrate buffer. Net photosynthesis rate was expressed as nmol of O₂.min⁻¹.cm⁻² of sediment surface.

Analyses on benthic invertebrates

Benthic invertebrates were collected by scraping a surface area of 150 cm² with a surber net in each sampled zone. Invertebrates were sorted and identified using Tachet et al. (2000) in the laboratory using a stereomicroscope. All taxa were classified into seven trophic functional groups (grazers, scrapers, deposit-feeders, filter-feeders, predators-parasites, detritivores and shredders) according to their trophic affinities (Tachet et al. 2000). When one taxon was affiliated to 2 or more functional groups, we applied a percentage of functional affinity in the abundance table for the given taxon (i.e., 50 % and 50 % for one taxon affiliated to 2 functional groups).

Statistical analyses

One-way ANOVAs were performed to compare biofilm variables and the abundances and richness (numbers of taxa) of each functional group of benthic invertebrates among wetlands. The relationship between the biofilm table and the fauna table was evaluated by a co-inertia analysis (see Dolédec and Chessel 1994; Dray et al. 2003). This multi-table analysis was performed to examine the links between the two tables, each table using the same 18 samples as rows. The relationship between each pair of data sets was measured using the Rv-coefficient (vectorial correlation coefficient; Robert and Escoufier 1976). The Rv-coefficient between two tables ranges between 0 and 1 and is a multidimensional equivalent of the ordinary correlation coefficient between two variables. For two given data sets X_k and X_l , the Rv-coefficient writes as follows:

$$Rv(X_k, X_l) = \frac{Covv(X_k, X_l)}{\sqrt{Vav(X_k)} \sqrt{Vav(X_l)}}$$

with $Covv(X_k, X_l) = \text{Trace} (X_k^t D_n X_l D_p)$ as the vectorial covariance and $Vav (X_k) = \text{Trace} (X_k^t D_n X_k D_p)$ and $Vav (X_l) = \text{Trace} (X_l^t D_n X_l D_q)$ as the vectorial variance for X_k and X_l ,

respectively; Matrix D_n contains row weights (common to both data sets) and matrices D_p and D_q contain the column weights of X_k and X_l , respectively (see Blanc et al. 1998). To test the significance of R_v -coefficients, rows were randomly permuted within tables. Permutations were repeated 1,000 times to obtain a null distribution of R_v -coefficients. We assessed the statistical significance by determining the proportion of null values that were greater than the observed R_v -coefficients.

Linear regressions were performed to determine the significance of the correlations among biofilm characteristics and the abundance and richness of invertebrate functional groups (e.g., green algae and grazer abundance). For all variables, the normality and the homoscedasticity of the residuals were verified using the Shapiro-Wilk's test and the Levene's test, respectively. ANOVA analyses, co-inertia analysis, linear regressions and graphical displays were performed using functions in the ade4 package (Chessel et al. 2004; Dray et al. 2007) in R (R Development Core team 2015). Significance for statistical tests was accepted for a first species risk $\alpha < 0.05$.

Results

Biofilm characteristics of the three wetlands

Algal colonization of the water-sediment interface was significantly different among wetlands (Fig. 2) and followed the gradient of net productivity measured (Fig. 1): green algae and diatom biomasses were positively correlated (Pearson's $r = 0.829$) and exhibited decreasing values from wetland A to wetland C with intermediate values in wetland B (one-way ANOVAs, $F_{2,15} = 58.9$, $p < 0.001$ for green algae and $F_{2,15} = 10.1$, $p < 0.01$ for diatoms). The densities of these two algal groups were positively correlated with the net photosynthetic activity ($R^2 = 0.65$, $p < 0.001$, $n = 18$

with green algae biomass, and $R^2 = 0.46$, $p < 0.01$, $n = 18$ with diatom biomass). Cyanobacteria biomass showed a distinct pattern as it was significantly lower in wetland B than in the two other wetlands (Fig. 2, one-way ANOVA, $F_{2,15} = 79.4$, $p < 0.001$). Percentages of total organic carbon and total nitrogen were also the lowest at the sediment interface of wetland B (one-way ANOVAs, $F_{2,15} = 5.1$, $p < 0.05$ for total organic carbon and $F_{2,15} = 5.9$, $p < 0.05$ for total nitrogen). Hence, we detected a significant and positive correlation between cyanobacteria biomass and total nitrogen ($R^2 = 0.37$, $p < 0.01$, $n = 18$) but not between cyanobacteria biomass and total organic carbon ($R^2 = 0.18$, $p > 0.05$, $n = 18$). In contrast, bacterial abundances were negatively correlated with cyanobacteria biomasses ($R^2 = 0.55$, $p < 0.001$, $n = 18$) and exhibited higher cell numbers in wetland B than in the other wetlands (one-way ANOVA, $F_{2,15} = 14.6$, $p < 0.001$).

Functional composition of benthic invertebrate communities in the three wetlands

Abundances and richness of grazers and scrapers were significantly different among wetlands (one-way ANOVAs, $F_{2,15} = 14.1$, $p < 0.001$ for grazer abundance, $F_{2,15} = 11.8$, $p < 0.001$ for scraper abundance, $F_{2,15} = 5.0$, $p < 0.05$ for grazer richness, and $F_{2,15} = 15.3$, $p < 0.001$ for scraper richness). Following the decreasing gradient of net productivity, they were the highest in wetland A and the lowest in wetland C with intermediate values in wetland B (Fig. 3). The abundance of filter-feeders tended to follow the same pattern with highest values in wetland A (Fig. 3, one-way ANOVA, $F_{2,15} = 7.8$, $p < 0.01$) although filter-feeder richness did not vary significantly among wetlands (one-way ANOVA, $F_{2,15} = 1.7$, $p > 0.22$). Deposit-feeders exhibited a particular distribution compared with grazers, scrapers and filter-feeders (Fig. 3). They were the most numerous in wetland B (one-way ANOVA, $F_{2,15} = 10.8$, $p < 0.01$) but their richness was low (between 1 and 2 taxa) and not

significantly different among wetlands (one-way ANOVA, $F_{2,15} = 2.5$, $p > 0.1$). In contrast with the other functional groups, the abundance and richness of predators were not significantly different among wetlands (Fig. 3, one-way ANOVA, $F_{2,15} = 2.4$, $p > 0.1$ for predator abundance and $F_{2,15} = 2.3$, $p > 0.15$ for predator richness).

Relationships between biofilm characteristics and functional composition of benthic invertebrate communities

The first two axis of co-inertia analysis explained more than 95% of the co-variation between biofilm and fauna datasets and the structure shared by the two datasets was significantly different from a structure expected from random association ($R_v = 0.60$, $p < 0.001$). Samples were discriminated among wetlands (Fig. 4A) and the short arrows displayed on Fig. 4A underlined the good fit between biofilm and fauna datasets in the shared structure.

On the factorial map of variables (Fig. 4BC), the horizontal axis indicated that samples characterized by high net photosynthetic activity and high diatom and green algae biomasses were also dense and rich in herbivores (grazers and scrapers). There were significant positive linear correlations between these biofilm characteristics (net photosynthetic activity, diatom or green algae biomass) and the densities of herbivores (grazers and scrapers) (Fig. 5). The richness of grazers and scrapers were also linearly and positively correlated with these biofilm characteristics and more especially with the biomass of green algae (relationships illustrated on Fig. 6). The vertical axis of the co-inertia analysis discriminated the samples with high bacterial densities and high abundances of deposit feeders from those of high cyanobacteria biomass and nitrogen content and low deposit-feeder abundances (Fig. 4BC). The abundance of deposit feeders was significantly and positively correlated with bacterial abundance (Fig. 7A) but not with total nitrogen and total

organic carbon ($R^2 < 0.1$, $p > 0.05$, $n = 18$). The abundance of deposit feeders was negatively correlated with cyanobacteria biomass (Fig. 7B). We did not detect clear correlations between biofilm characteristics and the functional groups of filter feeders and predators. The abundance of predators was significantly and positively correlated with the abundance of other invertebrates (Fig. 8). No significant correlations were obtained between predator richness and the abundance and richness of other invertebrates.

Discussion

Although the literature highlighted the occurrence of significant top-down control of communities in wetlands (Magnusson and Williams 2009; Hillebrand 2009; Moore 2018), top-down forces were apparently weak in our systems. Then, our dataset did not corroborate the prediction of EEH (Oksanen et al. 1981) suggesting a significant influence of top-down forces in our most productive wetland. The analysis of the links between biofilms and functional trophic groups of invertebrates clearly showed a dominant bottom-up effect of primary productivity (net photosynthesis) and algal resources on herbivores in artificial wetlands. Indeed, the abundances of both grazers and scrapers were positively and linearly correlated with the biomass of green algae and diatoms in wetlands. These correlations indicate that the bottom-up control of green algae and diatoms swamped out the top-down influence of herbivores on algal resources. As previous enclosure experiments showed that the introduction of a grazer (the gastropod *Viviparus viviparus*, a species which is not naturally present in our artificial wetlands) had a significant top-down effect on algae in these artificial wetlands (Gette-Bouvarot et al. 2015; Estragnat et al. 2018), we can conclude that pressure by

grazers and scrapers naturally present in these systems was not sufficient enough to reduce algal biomass.

Positive relationships between herbivore abundance and biomass of green algae and diatoms contrasted with the lack of relationship detected between herbivores and cyanobacteria biomass. This result may suggest that herbivores consumed green algae and diatoms rather than cyanobacteria in wetlands, resulting in no relationship between cyanobacteria biomass and herbivore abundance. This suggestion of a poor trophic transfer from cyanobacteria to herbivores is consistent with several studies indicating that cyanobacteria are generally non-attractive trophic sources for invertebrates due to their poor nutritional quality (low essential fatty acids and P contents, Gulati and DeMott 1997; Paerl and Fulton 2006) and their chemical and structural defense against grazing (DeMott et al. 1991; Soares et al. 2010; Bownik 2016). Our correlation analyses also suggest that cyanobacteria had a negative influence on deposit-feeders. However, a toxic impact of cyanobacteria on this functional group is highly improbable because examples of negative interactions between cyanobacteria and deposit feeders in the field are lacking in the literature and, in contrast, a study reported a positive effect of the fixation of N₂ by cyanobacteria on deposit feeders (Karlson et al. 2014). Moreover, we did not detect any negative influence of cyanobacteria biomass on grazers and scrapers, two trophic functional groups that would be the most affected by toxin produced by cyanobacteria (Bownik 2016). Together, these results suggest that cyanobacteria were probably not toxic for invertebrates in our artificial wetlands.

In addition to the positive relationship between algal (green algae and diatoms) biomass and herbivore abundances, we also showed that grazer and scraper richness were positively correlated with the biomasses of green algae, diatoms and photosynthetic activity. These results indicate that basal resources and net primary productivity may positively influence the horizontal diversity of the food web (*sensu* Srivastava and Bell 2009). This positive relationship between primary

productivity (and energy associated with trophic resources) and herbivore taxa richness can be related to the “species-energy theory” (Wright 1983; Srivastava and Lawton 1998; Evans et al. 2005). Indeed, several studies support the hypothesis that energy correlates positively with species richness (e.g, Huston 1994; Evans et al. 2005). In our field conditions, we can expect that an increase in benthic primary productivity could have elevated the amount of certain individual resource types (i.e., algal species), enabling specialist herbivore species to feed on them. Consequently, this trophic specialization would have led to reduced niche overlap among herbivores, reducing rates of competitive exclusion and elevating species richness in more productive wetlands. We also suspect that the diversity of individual resource types (i.e., algal species) increased with benthic primary productivity, enabling the highest herbivore diversity in the wetland characterized by the highest algal richness. Nevertheless, we cannot confirm this potential link between the trophic specialization of herbivore taxa and the abundances of individual resource types because green algae and diatom groups contained several species with contrasted trophic preferences for invertebrates (e.g., Tall et al. 2006; Alfaro 2009). More precise analyses such as invertebrate gut analyses using metabarcoding approaches (Pompanon et al. 2012) would be necessary to determine the trophic niche of each herbivore taxa in association with the availability of algal species in artificial wetlands. This perspective is very promising to evaluate whether the trophic specialization of herbivores in the most productive wetlands can explain the positive relationship between net benthic primary productivity (energy) and herbivore taxa richness in our artificial systems.

The positive and significant linear correlation between the abundances of predators and other aquatic invertebrates also suggest that bottom-up control of predators by their prey blurred the top-down control exerted by predators on their prey in our artificial wetlands. The lack of a strong top-down control of invertebrate predators on other invertebrates in the most productive wetland

(wetland A) would be associated with the increased horizontal diversity in food web observed for grazers and scrapers in this wetland. Indeed, increased diversity of a trophic level would mitigate top-down control by predators on this trophic level due to differences in predation pressure among the species of this trophic level. More precisely, a resource base with more species is more likely to contain at least one species that is relatively resistant to consumption and can compensate for the decrease of more consumable species in the presence of consumers (Leibold 1989, Duffy et al. 2007). However, the mechanism responsible of the lack of top-down cascade in our artificial wetlands is out of the scope of the present study. This further step would require more complete description and quantification of species-species interactions within the food webs.

Considering the detritus-based food web, the positive linear relationship between the bacterial abundance and the abundance of deposit-feeders suggests that the functional group of deposit-feeders was also bottom-up controlled by their trophic resources. Nevertheless, the abundances of bacteria and deposit-feeders were not correlated with the quantity of total organic carbon and total nitrogen in the sediment. These results are not consistent with the positive relationship between organic matter quantity and decomposer (micro- and macro-organisms) abundance classically supporting the bottom-up theory in sedimentary environments (Dannovaro et al. 1995; Lin and Yo 2008; Ramalho et al. 2014). The positive link between bacteria and deposit-feeder abundance observed in the present study was likely due to a positive influence of deposit-feeding activities on bacteria, a process largely demonstrated in a wide range of freshwater sediments (van de Bund et al. 1994; Griebler 1996; Traunspurger et al. 1997; Wieltschnig et al. 2008; Nogaro et al. 2009; Pignoret et al. 2016). A recent experiment performed in one of the artificial infiltration wetlands studied here demonstrated that aquatic oligochaetes, which are the main representatives of deposit-feeders in the present study, had a strongly positive and significant influence on the abundance and activity of bacteria in sediments (Mermillod-Blondin et al. 2018). More precisely, the bioturbation

associated with the addition of the tubificid worm *Limnodrilus hoffmeisteri* increased the bacterial abundance by 2-fold and the microbial hydrolytic activity by 35% without influencing the quantity of total nitrogen and organic carbon in sediments (Table 1). Thus, the top-down and positive effect of deposit-feeders on micro-organisms was the most probable mechanism explaining the positive relationship between bacteria and deposit-feeder abundances in the present experiment.

Conclusions

Our results did not corroborate the predictions of EEH suggesting that top-down controls would have a significant influence on community structure in the most productive ecosystems. Indeed, a dominance of bottom-up processes in the phototroph-based food web occurred in our artificial wetlands whatever their productivity. Our strong linear correlations between the biomasses and productivity of the algal component of the biofilm (especially green algae and diatoms) and the abundances of grazers and scrapers suggest that algal biomass and benthic primary productivity supported the herbivore trophic level. In contrast with the phototroph-based food web, results obtained for the detritus-based food web illustrate a positive effect of deposit-feeding invertebrates on bacteria. As observed in sedimentary environments with oligochaetes (Brinkhurst and Chua 1969; Wavre and Brinkhurst 1971; Fukuhara et al. 1980) and polychaetes (gardening effect, Hylleberg 1975), the feeding activities of deposit-feeder such as aquatic oligochaetes might have stimulated organic matter decomposition and bacterial growth in sediments. Thus, the present experiment showed that phototroph-based and detritus-based food webs were not dominated by the same processes in our artificial wetlands as herbivore abundance was likely controlled by the biomass of basal resources (algae) whereas deposit-feeder abundance was not linked to the amount of organic matter in sediments. Different feeding rates of invertebrates and contrasted renewal rate

of basal resources (more rapid for algae than for detritic organic matter) between the two food webs may explain the different links between invertebrates and basal food sources reported for both food webs. These links are expected to have implications on the whole ecosystem functioning (nutrient cycling, energy fluxes) and need to be considered in future studies. With this aim, development of labelling approaches using stable isotopes ^{13}C and ^{15}N - (Christianen et al. 2017) would be very promising to precisely quantify the transfer of energy within the phototrophic-based and detritus-based food webs and their consequences on ecosystem functioning. More research on artificial wetlands is needed because these artificial systems have been increasingly developed to compensate for the loss of natural wetlands, particularly in urbanized landscapes (Davidson 2014). This is a major challenge for the future, considering the dramatic increase of the urbanization and the role played by wetland ecosystems for biodiversity preservation and nutrient retention (Hanson et al. 2005).

Acknowledgments

We would like to thank Félix Vallier for his help during laboratory analyses. This research was done on the Research Platform of Crépieux-Charmy (Plate-forme de recherche de Crépieux-Charmy) and received financial and technical support from the Urban Community of Lyon (Grand Lyon La Métropole) and Veolia Water (Eau du Grand Lyon). This work was performed within the framework of the EUR H2O'Lyon (ANR-17-EURE-0018) of Université de Lyon (UdL), within the program "Investissements d'Avenir" operated by the French National Research Agency (ANR). We also would like to thank the two anonymous referees for their pertinent comments that significantly improved an earlier version of our manuscript.

404

405 References

406 Alfaro AC (2009) Diet of the pulmonate gastropod *Onchidella nigricans* in the intertidal rocky
407 shore, New Zealand. *Animal Biol* 59:231-240.

408 Aunapuu M, Dahlgren J, Oksanen T, Grellmann D, Oksanen L, Olofsson J, Rammul U, Schneider
409 M, Johansen B, Hygen HO (2007) Spatial patterns and dynamic responses of arctic food
410 webs corroborate the exploitation ecosystems hypothesis (EEH). *Am Nat* 171:249-262.

411 Blanc L, Chessel D, Dolédec S (1998) Etude de la stabilité temporelle des structures spatiales par
412 analyses d'une série de tableaux de relevés faunistiques totalement appariés. *Bull Fr Pêche*
413 *Pisc* 348:1-21.

414 Bertness MD, Brisson CP, Bevil MC, Crotty SM (2014) Herbivory drives the spread of salt marsh
415 die-off. *PLoS One* 9(3):e92916.

416 Bownik A (2016) Harmful algae: Effects of cyanobacterial cyclic peptides on aquatic invertebrates-
417 a short review. *Toxicon* 124:26-35.

418 Brinkhurst RO, Chua KE (1969) Preliminary investigation of the exploitation of some potential
419 nutritional resources by three sympatric tubificid Oligochaetes. *J Fish Res Bd Can* 26:2659-
420 68.

421 Chessel D, Dufour AB, Thioulouse J (2004) The ade4 package-I-One-table methods. *R news* 4: 5-
422 10.

423 Christianen MJA, Middelburg JJ, Holthuijsen SJ, Jouta J, Compton TJ, Heide T, Piersma T,
 424 Damsté JSS, van der Veer HW, Schouten S, Olff H (2017) Benthic primary producers are
 425 key to sustain the Wadden Sea food web: stable carbon isotope analysis at landscape scale.
 426 Ecology 98:1498-1512.

427 Cobbaert D, Bayley SE, Greter JL (2010) Effects of a top invertebrate predator (*Dytiscus*
 428 *alaskanus*; Coleoptera: Dytiscidae) on fishless pond ecosystems. *Hydrobiologia* 644(1):103-
 429 114.

430 Cummins KW, Klug MJ (1979) Feeding ecology of stream invertebrates. *Ann Rev Ecol Syst*
 431 10:147-172.

432 Danovaro R, Della Croce N, Eleftheriou A, Fabiano M, Papadopoulou N, Smith C, Tselepides A
 433 (1995) Meiofauna of the deep Eastern Mediterranean Sea: distribution and abundance in
 434 relation to bacterial biomass, organic matter composition and other environmental factors.
 435 *Prog Oceano* 36:329-341.

436 Davidson NC (2014) How much wetland has the world lost? Long-term and recent trends in global
 437 wetland area. *Mar. Freshwat Res* 65:934-941.

438 De Meester L, Declerck S, Stoks R, Louette G, Van De Meutter F, De Bie T, Michels E, Brendonck
 439 L (2005) Ponds and pools as model systems in conservation biology, ecology and
 440 evolutionary biology. *Aquatic conservation: Mar. Freshwat Ecosyst* 15:715-725.

441 DeMott WR, Zhang QX, Carmichael WW (1991) Effects of toxic cyanobacteria and purified toxins
 442 on the survival and feeding of a copepod and three species of *Daphnia*. *Limnol Oceano*
 443 36:1346-1357.

444 Diehl S (1992) Fish predation and benthic community structure: the role of omnivory and habitat
 445 complexity. *Ecology* 73:1646-1661.

446 Dolédec S, Chessel D (1994) Co-inertia analysis: an alternative method for studying species–
 447 environment relationships. *Freshwat Biol* 31:277-294.

448 Dray S, Chessel D, Thioulouse J (2003) Co-inertia analysis and the linking of ecological data
 449 tables. *Ecology* 84:3078-3089.

450 Dray S., Dufour AB, Chessel D (2007) The ade4 package-II: Two-table and K-table methods. *R*
 451 *news* 7:47-52.

452 Du X., García-Berthou E., Wang Q., Liu J., Zhang T., Li Z. (2015) Analyzing the importance of
 453 top-down and bottom-up controls in food webs of Chinese lakes through structural equation
 454 modeling. *Aquat Ecol* 49:199-210.

455 Duffy JE, Cardinale BJ, France KE, McIntyre PB, Thébault E, Loreau M (2007) The functional
 456 role of biodiversity in ecosystems: incorporating trophic complexity. *Ecology Lett* 10:522–
 457 538.

458 Estragnat V, Mermillod-Blondin F, Jully M; Lemoine D; Lassabatere L; Volatier L (2018) Does
 459 the efficiency of grazer introduction to restore and preserve the hydraulic performance of
 460 infiltration basins depend on the physical and biological characteristics of the infiltration
 461 media? *Ecol Engin* 116:127-132.

462 Evans KL, Warren PH, Gaston KJ (2005) Species–energy relationships at the macroecological
 463 scale: a review of the mechanisms. *Biol Rev* 80:1-25.

464 Foulquier A, Malard F, Mermillod-Blondin F, Montuelle B, Dolédec S, Volat B, Gibert J (2011)
 465 Surface water linkages regulate trophic interactions in a groundwater food web. *Ecosystems*
 466 14:1339-1353.

467 Fukuhara H, Kikuchi E, Kurihara Y (1980) The effects of *Branchiura sowerbyi* (Tubificidae) on
 468 bacterial populations in submerged ricefield soil. *Oikos* 34:88-93.

469 Gette-Bouvarot M, Mermillod-Blondin F, Angulo-Jaramillo R, Delolme C, Lemoine D,
 470 Lassabatere L, Loizeau S, Volatier L (2014) Coupling hydraulic and biological
 471 measurements highlights the key influence of algal biofilm on infiltration basin performance.
 472 *Ecohydrology* 7:950-964.

473 Gette-Bouvarot M, Volatier L, Lassabatere L, Lemoine D, Simon L, Delolme C, Mermillod-
 474 Blondin F (2015) Ecological engineering approaches to improve hydraulic properties of
 475 infiltration basins designed for groundwater recharge. *Env Sci Technol* 49: 9936-9944.

476 Griebler C (1996) Some applications for the DMSO-reduction method as a new tool to determine
 477 the microbial activity in water-saturated sediments. *Arch Hydrobiol Suppl* 113:405–410.

478 Gruner DS, Smith JE, Seabloom EW, Sandin SA, Ngai JT, Hillebrand H, Harpole WS, Elser JJ,
 479 Cleland EE, Bracken MES, Borer ET, Bolker BM (2008) A cross-system synthesis of
 480 consumer and nutrient resource control on producer biomass. *Ecol Lett* 11:740-755.

481 Gulati R, DeMott W (1997) The role of food quality for zooplankton: remarks on the state-of-the-
 482 art, perspectives and priorities. *Freshwat Biol* 38:753-768.

483 Hansson LA, Brönmark C, Anders Nilsson P, Åbjörnsson K (2005) Conflicting demands on
484 wetland ecosystem services: nutrient retention, biodiversity or both? *Freshwat Biol* 50:705-
485 714.

486 Harris T, Graham J (2015) Preliminary evaluation of an *in vivo* fluorometer to quantify algal
487 periphyton biomass and community composition. *Lake Reserv Manag* 31:127–133.

488 Hillebrand H (2009) Meta-analysis of grazer control of periphyton biomass across aquatic
489 ecosystems. *J Phycol* 45:798-806.

490 Hoset KS, Ruffino L, Tuomi M, Oksanen T, Oksanen L, Mäkynen A, Johansen B, Moe T (2017)
491 Changes in the spatial configuration and strength of trophic control across a productivity
492 gradient during a massive rodent outbreak. *Ecosystems* 20:1421-35.

493 Hunter MD, Price PW (1992) Playing chutes and ladders: heterogeneity and the relative roles of
494 bottom-up and top-down forces in natural communities. *Ecology* 73:724-732.

495 Huston MA (1994) Biological diversity: the coexistence of species. Cambridge University Press.

496 Hylleberg J (1975) Selective feeding by *Abarenicola pacifica* with notes on *Abarenicola*
497 *vagabunda* and a concept of gardening in lugworms. *Ophelia* 14:113-137.

498 Karlson AM, Gorokhova E, Elmgren R (2014) Nitrogen fixed by cyanobacteria is utilized by
499 deposit-feeders. *PLoS One* 9:e104460.

500 Laske SM, Rosenberger AE, Kane WJ, Wipfli MS, Zimmerman CE (2017) Top-down control of
501 invertebrates by Ninespine Stickleback in Arctic ponds. *Freshwat Sci* 36:124-137.

502 Leibold MA (1989) Resource edibility and the effects of predators and productivity on the outcome
503 of trophic interactions. *Am Nat* 134:922-949.

504 Leibold MA, Chase JM, Shurin JB, Downing AL (1997) Species turnover and the regulation of
505 trophic structure. *Annu Rev Ecol Syst* 28:467-494.

506 Lezama F, Baeza S, Altesor A, Cesa A, Chaneton EJ, Paruelo JM (2014) Variation of grazing-
507 induced vegetation changes across a large-scale productivity gradient. *J Veg Sci* 25:8-21.

508 Lin KJ, Yo SP (2008) The effect of organic pollution on the abundance and distribution of aquatic
509 oligochaetes in an urban water basin, Taiwan. *Hydrobiologia* 596:213-223.

510 Magnusson AK, Williams DD (2009) Top-down control by insect predators in an intermittent
511 pond—a field experiment. *Annal Limnol-Internat J Limnol* 45:131-143.

512 Menge BA (2000) Top-down and bottom-up community regulation in marine rocky intertidal
513 habitats. *J Exp Mar Biol Ecol* 250:257-289.

514 Menge BA, Olson AM, Dahlhoff EP (2002) Environmental stress, bottom-up effects, and
515 community dynamics: integrating molecular-physiological and ecological approaches. *Integr*
516 *Compar Biol* 42:892-908.

517 Mermillod-Blondin F, Simon L, Maazouzi C, Foulquier A, Delolme C, Marmonier P (2015)
518 Dynamics of dissolved organic carbon (DOC) through stormwater basins designed for
519 groundwater recharge in urban area: Assessment of retention efficiency. *Water Res* 81:27–
520 37.

521 Mermillod-Blondin F, Bouvarot M, Déjollat Y, Adrien J, Maire E, Lemoine D, Marmonier P,
522 Volatier L (2018) Influence of tubificid worms on sediment structure, benthic biofilm and
523 fauna in wetlands: A field enclosure experiment. *Freshwat Biol* 63:1420-1432.

524 Moore A (2018) Context-dependent consumer control in New England tidal wetlands. *PloS One*
525 13(5):e0197170.

526 Nogaro G, Mermillod-Blondin F, Valett MH, François-Carcaillet F, Gaudet J-P, Lafont M., Gibert
527 J (2009) Ecosystem engineering at the sediment-water interface: Bioturbation and consumer-
528 substrate interaction. *Oecologia* 161:125–138.

529 Odum EP, Smalley AE (1959) Comparison of population energy flow of an herbivorous and a
530 deposit-feeding invertebrate in a saltmarsh ecosystem. *Proc Natl Acad Sci USA* 45:617–622.

531 Oksanen L, Fretwell SD, Arruda J, Niemela P (1981) Exploitation ecosystems in gradients of
532 primary productivity. *Am. Nat.* 118:240-261.

533 Paerl HW, Fulton RS (2006) Ecology of harmful cyanobacteria. p. 95-109. In: E Granéli and JT
534 Turner [eds.], *Ecology of harmful algae*. Springer, Berlin, Heidelberg.

535 Pigneret M, Mermillod-Blondin F, Volatier L, Romestaing C, Maire E, Adrien J, Guillard L,
536 Roussel D, Hervant F (2016) Urban pollution of sediments: Impact on the physiology and
537 burrowing activity of tubificid worms and consequences on biogeochemical processes. *Sci*
538 *Tot Env* 568:196-207.

539 Polis GA, Strong DR (1996) Food web complexity and community dynamics. *Am Nat* 147:813-
540 846.

541 Pompanon F, Deagle BE, Symondson WO, Brown DS, Jarman SN, Taberlet P (2012) Who is eating
542 what: diet assessment using next generation sequencing. *Mol Ecol* 21:1931-50.

543 Ramalho SP, Adão H, Kiriakoulakis K, Wolff GA, Vanreusel A, Ingels J (2014) Temporal and
544 spatial variation in the Nazaré Canyon (Western Iberian margin): Inter-annual and canyon
545 heterogeneity effects on meiofauna biomass and diversity. *Deep Sea Res Part I: Oceano Res*
546 *Papers* 83:102-114.

547 Robert P, Escoufier Y (1976) A unifying tool for linear multivariate statistical methods: the RV-
548 coefficient. *Appl Stat* 25:257-265.

549 Sala NM, Bertness MD, Silliman B (2008) The dynamics of bottom-up and top-down control in a
550 New England salt marsh. *Oikos* 117:1050-1056.

551 Shurin JB, Borer ET, Seabloom EW, Anderson K, Blanchette CA, Broitman B, Cooper SD,
552 Halpern BS (2002) A cross-ecosystem comparison of the strength of trophic cascades. *Ecol*
553 *Lett* 5:785-791.

554 Soares MCS, Lürling M, Huszar VL (2010) Responses of the rotifer *Brachionus calyciflorus* to
555 two tropical toxic cyanobacteria (*Cylindrospermopsis raciborskii* and *Microcystis*
556 *aeruginosa*) in pure and mixed diets with green algae. *J Plankton Res* 32:999-1008.

557 Srivastava DS, Bell T (2009) Reducing horizontal and vertical diversity in a foodweb triggers
558 extinctions and impacts functions. *Ecol Lett* 12:1016-1028.

559 Srivastava DS, Lawton JH (1998) Why more productive sites have more species: an experimental
560 test of theory using tree-hole communities. *Am Nat* 152:510-529.

- 561 Steiner CF (2001) The effects of prey heterogeneity and consumer identity on the limitation of
562 trophic-level biomass. *Ecology* 82:2495-2506.
- 563 Sundbäck K, Miles A, Hulth S, Pihl L, Engström P, Selander E, Svenson A (2003) Importance of
564 benthic nutrient regeneration during initiation of macroalgal blooms in shallow bays. *Mar*
565 *Ecol Prog Ser* 246:115-126.
- 566 Tachet H, Richoux P, Bournaud M, Usseglio-Polatera P (2000) Invertébrés d'eau douce:
567 systématique, biologie, écologie. CNRS éditions, Paris.
- 568 Tall L., Cattaneo A., Cloutier L., Dray S., Legendre P (2006) Resource partitioning in a grazer
569 guild feeding on a multilayer diatom mat. *J North Am Benthol Soc* 25:800-810.
- 570 Traunspurger W, Bergtold M, Goedkoop W (1997) The effects of nematodes on bacterial activity
571 and abundance in a freshwater sediment. *Oecologia* 112:118-122.
- 572 Van de Bund WJ, Goedkoop W, Johnson RK (1994) Effects of deposit-feeder activity on bacterial
573 production and abundance in profundal Lake sediment. *J North Am Benthol Soc* 13:532-539.
- 574 Van de Koppel J, Herman PM, Thoolen P, Heip CH (2001) Do alternate stable states occur in
575 natural ecosystems? Evidence from a tidal flat. *Ecology* 82:3449-3461.
- 576 Watermann F, Hillebrand H, Gerdes G, Krumbein WE, Sommer U (1999) Competition between
577 benthic cyanobacteria and diatoms as influenced by different grain sizes and temperatures.
578 *Mar Ecol Prog Ser* 187:77-87.
- 579 Wavre M, Brinkhurst RO (1971) Interactions between some tubificid oligochaetes and bacteria
580 found in the sediments of Toronto Harbour, Ontario. *J Fish Res Bd Can* 28:335-341.

581 Wieltschnig C, Fischer UR, Velimirov B, Kirschner AK (2008) Effects of deposit-feeding
582 macrofauna on benthic bacteria, viruses, and protozoa in a silty freshwater sediment. *Microb*
583 *Ecol* 56:1-12.

584 Wright DH (1983) Species-energy theory: an extension of species-area theory. *Oikos* 41:496-506.

585

Figure legends:

Fig 1: Percentage of fine particles ($< 63 \mu\text{m}$), benthic algal biomass and net photosynthetic activity of benthic biofilms measured in the three studied wetlands ($n = 6$ measures per wetland).

Fig 2: Characteristics of the biofilms in the three wetlands ($n = 6$ measures per wetland for each variable).

Fig 3: Abundances (ind/m^2) and richness (taxa/sample) of the five major trophic functional groups of benthic invertebrates collected in the three wetlands ($n = 6$ measures per wetland).

Fig 4: Co-inertia analysis showing a) the shared structure obtained from the biofilm and fauna datasets, b) the factorial map of biofilm variables and c) the factorial map of variables associated with invertebrate functional groups. A = Wetland A, B = Wetland B, C = Wetland C, PhotoS = Net Photosynthesis rate, TOC = Total organic carbon, TN = Total nitrogen, DepositAB = Abundances of deposit feeders, DepositRICH = Richness of deposit feeders, ScrapAB = Abundances of scrapers, ScrapRICH = Richness of scrapers, FiltAB = Abundances of filter-feeders, FiltRICH = Richness of filter-feeders, GrazerAB = Abundances of grazers, GrazerRICH = Richness of grazers, PredAB = Abundances of predators, PredRICH = Richness of predators.

Fig 5: Correlations a) between net photosynthetic activity and grazer abundance, b) between net photosynthetic activity and scraper abundance, c) between green algae biomass and grazer abundance, d) between green algae biomass and scraper abundance, e) between diatom biomass and grazer abundance, and f) between diatom biomass and scraper abundance.

Fig 6: Correlations a) between green algae biomass and grazer richness and b) between green algae biomass and scraper richness.

607 Fig 7: Correlations a) between bacterial abundance and deposit-feeder abundance, and b) between
608 cyanobacteria biomass and deposit-feeder abundance.

609 Fig 8: Correlation between predator abundance and the abundance of other invertebrates.

610

611

612

613

614

Fig 1

615

Fig 2

Fig 3

Fig 4

Fig 5

623

624

Fig 6

Fig 7

Fig 8

629

630 Table 1. Mean values ($\pm$ SD) of the biofilm variables measured in 6 enclosures without worms and
 631 6 enclosures in which a density of 20,000 individuals.m⁻² of the tubificid worm *L. hoffmeisteri* has
 632 been added. Student-t tests were run to evaluate the influence of tubificid worm addition on biofilm
 633 variables. Modified from Mermillod-Blondin et al. (2018)
 634
 635

Measured variables	Control enclosures averaged values $\pm$ standard deviation	Worm enclosures averaged values $\pm$ standard deviation	t _(10, 0.05)	p-values
Bacterial cells (10 ⁹ /g of dry sediment)	3.20 $\pm$ 1.05	6.48 $\pm$ 1.90	4.04	0.002
Total organic carbon (% of dry sediment)	0.788 $\pm$ 0.424	0.704 $\pm$ 0.268	0.41	0.689
Total nitrogen (% of dry sediment)	0.117 $\pm$ 0.056	0.112 $\pm$ 0.034	0.19	0.857
Hydrolytic activity (10 ⁻² μ mol of hydrolyzed FDA .h ⁻¹ .g ⁻¹ of dry sediment)	4.57 $\pm$ 1.00	6.16 $\pm$ 0.84	2.99	0.013

636

637

638 Supplementary Material 1: Table of taxa abundances for each sample (sampled surface: 150 cm²).
639 The last column indicates the affiliation of taxa to functional groups. G: Grazers, S: Scrapers, DF:
640 Deposit-feeders, F: filter-feeders, P: predators and parasites, D: Detritivores, and Shr: Shredders.

641

Taxa	Sampled basin						Feeding groups
	A	A	A	A	A	A	
<i>Naidinae sp.</i>	35	32	47	57	56	104	G/S
other oligochaetes	30	13	7	8	8	8	G/S
Enchytraeidae	116	82	127	235	464	104	DF
<i>Hydra sp.</i>	54	6	32	27	24	152	P
<i>Dugesia spp.</i>	4	0	0	1	8	0	P
Nematoda	98	63	58	36	568	168	P/DF
<i>Erpobdella sp.</i>	0	0	0	0	0	0	P
<i>Physa acuta</i> + <i>Physa fontinalis</i>	7	5	2	4	8	0	G/S
<i>Galba truncatula</i>	161	45	44	40	168	88	G/S
<i>Gyraulus sp.</i>	0	0	0	2	0	8	G/S
Hydracarina	0	0	0	1	0	0	P
Cyclopoida	721	277	740	524	1048	1624	F
Calanoida	0	0	0	0	0	0	F
Harpacticoida	0	2	0	0	0	0	B
<i>Chidorus sphaericus</i> + <i>Chidorus sp.</i>	14	0	120	0	145	94	F
<i>Eurycercus lamellatus</i>	95	40	150	137	181	188	F
<i>Sida crystallina</i>	27	26	180	0	0	141	F
<i>Simocephalus vetulus</i>	191	40	120	0	36	423	F
<i>Simocephalus expinosus</i>	109	92	209	137	508	470	F
<i>Simocephalus sp.</i>	0	13	60	55	36	0	F
<i>Alona quadrangularis</i>	82	0	30	0	109	141	F
<i>Pleuroxus sp.</i>	14	0	0	0	36	0	F
<i>Acroperus harpae</i>	0	13	0	0	36	0	F
<i>Daphnia pulex</i>	123	357	568	906	290	800	F
Non identified cladocera + other Daphnia	14	0	60	0	109	94	F
<i>Pseudocandona albicans</i>	0	0	0	1	0	0	DF
<i>Ilyocypris bradyi</i>	0	0	0	1	0	0	DF
<i>Heterocypris salina</i>	4	2	0	4	0	0	DF

642

643

644

645 Suppl Material 1 (continued)

646

Taxa	Sampled basin						Feeding groups
	A	A	A	A	A	A	
<i>Cypridopsis vidua</i>	20	1	0	9	0	8	DF
<i>Bradleystrandesia obliqua</i>	22	6	8	7	16	0	F
<i>Limmocythere inopinata</i>	0	0	0	0	0	0	DF
<i>Gammarus sp.</i>	1	0	0	0	0	0	Shr/P
<i>Cloeon sp.</i>	43	76	95	152	208	224	B
<i>Caenis sp.</i>	0	0	0	0	0	0	Shr/D
<i>Hydroptilidae agrayla</i>	11	3	0	3	2	4	G/S
<i>Haliplidae haliplus</i>	6	6	0	4	1	0	Shr
<i>Gyrinidae gyrinus</i> + <i>Hydrophilidae berosus</i> + <i>Elmidae esolus</i>	1	1	0	0	0	0	P
Chironomidae Tanytarsini	145	108	116	48	264	112	F/DF
Chironomidae Chironominae	85	31	41	6	48	56	F/DF
Chironomidae Orthocladinae	58	18	33	50	32	88	R/DF
Chironomidae Tanypodinae	11	7	9	0	0	8	P
Nymphs of chironomidae	44	8	43	29	24	16	P/F/DF/S
<i>Chaoborus sp.</i>	1	0	0	0	0	0	P
Ceratopogoninae	2	0	0	3	0	0	D
Dasyhelinae	0	0	0	3	1	3	D
Other diptera	3	0	0	6	1	3	P
Libellulidae + Sympetrinae + Plactycnemididae	5	0	1	4	2	0	P

647

648

649

650

651 Suppl Material 1 (continued)

652

Taxa	Sampled basin						Feeding groups
	B	B	B	B	B	B	
<i>Naidinae sp.</i>	64	40	87	53	60	23	G/S
other oligochaetes	32	0	1	0	10	3	G/S
Enchytraeidae	640	1000	1289	699	1191	415	DF
<i>Hydra sp.</i>	0	0	1	0	4	0	P
<i>Dugesia spp.</i>	0	0	4	0	0	0	P
Nematoda	128	128	50	102	104	18	P/DF
<i>Erpobdella sp.</i>	0	0	0	0	0	0	P
<i>Physa acuta</i> + <i>Physa fontinalis</i>	8	0	2	1	2	0	G/S
<i>Galba truncatula</i>	160	64	43	131	104	40	G/S
<i>Gyraulus sp.</i>	0	0	0	0	0	0	G/S
Hydracarina	0	0	0	0	0	1	P
Cyclopoida	312	248	238	219	200	68	F
Calanoida	0	0	0	0	0	0	F
Harpacticoida	0	0	2	2	0	0	B
<i>Chidorus sphaericus</i> + <i>Chidorus sp.</i>	38	25	0	14	48	7	F
<i>Eurycercus lamellatus</i>	19	0	39	12	12	3	F
<i>Sida crystallina</i>	19	25	65	16	83	13	F
<i>Simocephalus vetulus</i>	0	0	45	16	24	6	F
<i>Simocephalus expinosus</i>	29	62	39	0	12	14	F
<i>Simocephalus sp.</i>	0	0	0	0	0	1	F
<i>Alona quadrangularis</i>	19	25	0	21	24	10	F
<i>Pleuroxus sp.</i>	0	0	0	0	0	2	F
<i>Acroperus harpae</i>	29	0	0	5	0	0	F
<i>Daphnia pulex</i>	0	12	13	21	0	0	F
Non identified cladocera + other Daphnia	0	12	0	11	12	0	F
<i>Pseudocandona albicans</i>	0	0	11	2	1	0	DF
<i>Ilyocypris bradyi</i>	0	8	0	3	2	0	DF
<i>Heterocypris salina</i>	0	0	0	0	0	0	DF

653

654

655

656

657 Suppl Material 1 (continued)
658

Taxa	Sampled basin						Feeding groups
	B	B	B	B	B	B	
<i>Cypridopsis vidua</i>	0	24	6	11	5	0	DF
<i>Bradleystrandesia obliqua</i>	37	16	9	14	9	0	F
<i>Limmocythere inopinata</i>	19	0	2	0	0	0	DF
<i>Gammarus sp.</i>	0	0	0	0	0	0	Shr/P
<i>Cloeon sp.</i>	32	24	34	18	55	19	B
<i>Caenis sp.</i>	0	0	3	1	0	0	Shr/D
<i>Hydroptilidae agrayla</i>	0	0	0	3	1	1	G/S
<i>Haliplidae haliplus</i>	4	0	4	7	9	1	Shr
<i>Gyrinidae gyrinus + Hydrophilidae berosus + Elmidae esol</i>	1	0	0	0	0	0	P
Chironomidae Tanytarsini	464	243	210	292	177	65	F/DF
Chironomidae Chironominae	137	46	131	171	135	56	F/DF
Chironomidae Orthocladinae	61	21	15	35	15	7	R/DF
Chironomidae Tanypodinae	15	5	3	14	4	7	P
Nymphs of chironomidae	61	5	15	13	16	3	P/F/DF/S
<i>Chaoborus sp.</i>	0	0	0	0	0	0	P
Ceratopogoninae	0	0	0	1	0	0	D
Dasyhelinae	0	0	0	0	0	0	D
Other diptera	1	0	0	1	0	0	P
Libellulidae + Sympetrinae + Plactycnemididae	0	0	3	2	1	0	P

659

660

661

662

663 Suppl Material 1 (continued)

664

			Sampled basin				
Taxa	C	C	C	C	C	C	Feeding groups
<i>Naidinae sp.</i>	4	1	1	0	0	0	G/S
other oligochaetes	0	3	0	0	8	0	G/S
Enchytraeidae	16	13	101	28	432	724	DF
<i>Hydra sp.</i>	1	0	0	2	0	0	P
<i>Dugesia spp.</i>	0	0	0	0	0	0	P
Nematoda	149	115	198	202	152	331	P/DF
<i>Erpobdella sp.</i>	0	1	0	1	0	0	P
<i>Physa acuta</i> + <i>Physa fontinalis</i>	4	3	1	8	0	0	G/S
<i>Galba truncatula</i>	19	14	17	7	32	37	G/S
<i>Gyraulus sp.</i>	1	0	0	0	0	0	G/S
Hydracarina	0	1	1	0	0	1	P
Cyclopoida	115	42	48	168	96	266	F
Calanoida	28	15	71	76	272	208	F
Harpacticoida	0	0	0	1	0	0	B
<i>Chidorus sphaericus</i> + <i>Chidorus sp.</i>	19	0	0	16	0	100	F
<i>Eurycercus lamellatus</i>	0	0	19	62	56	166	F
<i>Sida crystallina</i>	9	0	19	31	0	66	F
<i>Simocephalus vetulus</i>	0	0	10	0	0	33	F
<i>Simocephalus expinosus</i>	19	0	38	47	0	166	F
<i>Simocephalus sp.</i>	0	0	0	0	0	0	F
<i>Alona quadrangularis</i>	9	0	0	0	0	33	F
<i>Pleuroxus sp.</i>	37	0	0	0	0	0	F
<i>Acroperus harpae</i>	0	0	0	0	0	0	F
<i>Daphnia pulex</i>	346	372	392	622	1324	763	F
Non identified cladocera + other Daphnia	19	16	0	0	28	166	F
<i>Pseudocandona albicans</i>	1	0	8	8	0	0	DF
<i>Ilyocypris bradyi</i>	0	0	0	0	0	0	DF
<i>Heterocypris salina</i>	0	0	0	0	0	0	DF

665

666

667

668

669 Suppl Material 1 (continued)

670

Taxa	Sampled basin						Feeding groups
	C	C	C	C	C	C	
<i>Cypridopsis vidua</i>	2	0	4	8	0	1	DF
<i>Bradleystrandesia obliqua</i>	3	3	12	40	0	2	F
<i>Limmocythere inopinata</i>	0	0	0	0	0	0	DF
<i>Gammarus sp.</i>	0	1	0	0	0	0	Shr/P
<i>Cloeon sp.</i>	16	7	27	54	27	26	B
<i>Caenis sp.</i>	0	0	0	0	0	0	Shr/D
<i>Hydroptilidae agrayla</i>	0	0	1	0	0	0	G/S
<i>Haliplidae haliplus</i>	0	0	0	0	0	0	Shr
<i>Gyrinidae gyrinus + Hydrophilidae berosus + Elmidae esol</i>	0	0	2	0	1	1	P
Chironomidae Tanytarsini	40	6	51	31	16	121	F/DF
Chironomidae Chironominae	16	2	13	7	8	2	F/DF
Chironomidae Orthocladinae	0	0	2	1	0	0	R/DF
Chironomidae Tanypodinae	5	1	3	2	0	1	P
Nymphs of chironomidae	2	1	0	2	8	1	P/F/DF/S
<i>Chaoborus sp.</i>	0	0	2	1	0	1	P
Ceratopogoninae	0	0	0	1	0	0	D
Dasyhelinae	0	0	0	0	0	0	D
Other diptera	0	0	2	2	0	1	P
Libellulidae + Sympetrinae + Plactycnemididae	1	2	1	2	0	0	P

671

672

673

674

675

676